

**Marked-Up Copy of Claims, as Amended  
In the Amendment Responding to the  
Office Action Dated 17 July 2001**

2. (Amended) A method of identifying a gene, expression of which is regulated by hydrodynamic stress, the method comprising

a) subjecting a first vascular endothelial cell of a mammal to hydrodynamic stress~~The method of claim 1, wherein the cell is subjected to hydrodynamic stress using an in vitro flow chamber;~~

b) thereafter assessing the level of expression of the gene in the cell; and

c) comparing the level of expression of the gene in the first cell with the level of expression of the same gene in a second vascular endothelial cell of the mammal, the second cell being subjected to different hydrodynamic stress;

whereby a difference between the level of expression of the gene in the first cell and the level of expression of the gene in the second cell is an indication that the gene is regulated by hydrodynamic stress.

5. (Amended) The method of claim-1 2, wherein the mammal is a human.

6. (Amended) The method of claim-1 2, wherein the hydrodynamic stress is exerted in an amount from about 0 dyn/cm<sup>2</sup> to about 100 dyn/cm<sup>2</sup>.

7. (Amended) The method of claim-1 2, wherein the hydrodynamic stress is exerted for at least about several seconds.

8. (Amended) The method of claim ~~1~~ 2, wherein the cell is a single cell isolated in vivo.

10. (Amended) The method of claim ~~1~~ 2, wherein ~~the cell is~~ a group of cells is isolated in vivo.

11. (Amended) The method of claim ~~1~~ 2, wherein the cell is a single cell isolated in vitro from a confluent monolayer.

12. (Amended) The method of claim ~~1~~ 2, wherein the cell is one of a group of cells isolated in vitro from a confluent monolayer.

13. (Amended) The method of claim ~~1~~ 2, wherein the level of expression of the gene is assessed using amplified antisense RNA in combination with northern blotting or a microarray technique.

19. (Amended) ~~The method of claim 18, wherein the cell is subjected to hydrodynamic stress~~ A method of identifying a nucleic acid comprising a hydrodynamic stress regulation (HSR) region, the method comprising

a) subjecting a first vascular endothelial cell of a mammal to hydrodynamic stress using an in vitro flow chamber capable of generating a hydrodynamic stress flow field having spatially defined microheterogeneity;

b) thereafter assessing the level of expression of the nucleic acid in the cell; and

c) comparing the level of expression of the nucleic acid in the first cell with the level of expression of the same nucleic acid in a second vascular endothelial cell of the mammal, the second cell being subjected to different hydrodynamic stress;

whereby a difference between the level of expression of the nucleic acid in the first cell and the level of expression of the nucleic acid in the second cell is an indication that the nucleic acid comprises a HSR region.

20. (Amended) The method of claim ~~18~~ 19, wherein the level of expression of the nucleic acid is assessed using amplified antisense RNA from a single endothelial cell.

21. (Amended) A method of identifying a nucleic acid comprising an HSR region, the method comprising

a) comparing the sequence of a first nucleic acid with the sequence of a second nucleic acid comprising an HSR region, the second nucleic acid being identified by the method of claim 19; and

b) identifying a region of the first nucleic acid which is homologous to the HSR region of the second nucleic acid, whereby a nucleic acid comprising a HSR region is identified.

23. (Amended) ~~The method of claim 22, wherein the cell is subjected~~ A method of identifying a hydrodynamic stress responsive protein, the method comprising

a) subjecting a first vascular endothelial cell of a mammal to hydrodynamic stress using an in vitro flow chamber capable of generating a hydrodynamic stress flow field having spatially defined microheterogeneity;

b) thereafter assessing the level of expression of a nucleic acid in the first cell;

c) comparing the level of expression of the nucleic acid in the first cell with the level of expression of the same nucleic acid in a second vascular endothelial cell of the mammal, the second cell being subjected to different hydrodynamic stress, whereby a difference between the level of expression of the nucleic acid in the first cell and the level of expression of the nucleic acid in the second cell is an indication that the nucleic acid encodes a hydrodynamic stress responsive protein; and

d) identifying a protein encoded by the nucleic acid, whereby a hydrodynamic stress responsive protein is identified.

24. (Amended) The method of claim-~~22~~ 23, wherein the level of expression of the nucleic acid is assessed in a single endothelial cell.

25. (Amended) ~~An~~ A method of making an array of nucleic acids comprising an HSR region, the method comprising

a) selecting a plurality of nucleic acids comprising an HSR region wherein at least one of the nucleic acids is selected by the method of claim-~~18~~ 19; and

b) incorporating the selected nucleic acids into an array.

26. (Amended) A kit for carrying out the method of claim-~~1~~ 2, the kit comprising

- a) an instructional material describing use of the kit for the method of claim 2;
- b) a reagent for use in amplified antisense RNA;
- c) a reagent for use in northern blotting or microarray analysis; and
- d) a radiolabeled or fluorescent probe.